

POSTER PRESENTATION

Open Access

MAP kinase inhibitors stimulate T cell and anti-tumor activity in combination with blockade of the PD-L1/PD-1 interaction

Bryan A Irving*, Jeanne Cheung, Yagai Yang, Marina Moskalenka, Marcin Kowanetz, Heather Maecker, Ira Mellman

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Pharmacological inhibition of the MAPK pathway with MEK or BRAF antagonists has proved successful in inducing regression of melanoma tumors bearing the targeted activating mutations. Moreover, antibodies targeting T-cell immune checkpoint inhibitors CTLA-4 or PD-L1/PD-1 have demonstrated the capacity to generate durable responses in patients with multiple cancer types. Thus, combining MAPK pathway-targeted agents with antibodies that enhance anti-tumor immunity represents an increasingly attractive treatment paradigm for cancer. However, little is known about the impact of tumor-targeted agents on immune function as similar signaling pathways drive both T-cell activation and cancer cell proliferation. Accordingly, agents targeting MAPK-dependent tumor growth would be predicted to also inhibit T-cell immunity. Here we show that, unexpectedly, potent suppression of T-cell receptor (TCR) function by MEK inhibition can be largely overcome in the presence of co-stimulation by anti-CD28 in vitro or blockade of the inhibitory PD-L1/PD-1 pathway in T cells in vivo. The ability of anti-CD28 to override suppression of T-cell activation by MEK inhibitors was dependent on the PI3K/mTOR pathway. Enhanced anti-tumor activity was also observed combining MEK inhibition with PD-L1 blockade, which was likely potentiated by upregulation of tumor MHC Class I expression through inhibition of MEK. Interestingly, inhibitors targeting BRAF V600E mutations actually augmented TCR-driven proliferation in vitro and T-cell function in vivo when combined with a vaccine or blockade of PD-L1 exclusively in the context of a wildtype BRAF background. These data demonstrate that targeting the MAPK pathway can be compatible with or even

enhance T-cell function and provide rationale for combining these inhibitors with immunotherapy in clinical trials.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P79

Cite this article as: Irving et al.: MAP kinase inhibitors stimulate T cell and anti-tumor activity in combination with blockade of the PD-L1/PD-1 interaction. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P79.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Genentech, Inc., South San Francisco, CA, USA



© 2013 Irving et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.